

On the Classicality of Broda's $SU(2)$ Invariants of 4-Manifolds

Louis Crane¹

*Department of Mathematics
Kansas State University
Manhattan, KS 66506-2602*

Louis H. Kauffman²

*Department of Mathematics, Statistics and Computer Science
University of Illinois at Chicago
Chicago, IL 60680*

David N. Yetter

*Department of Mathematics
Kansas State University
Manhattan, KS 66506-2602*

Abstract: Recent work of Roberts [R] has shown that the surgical 4-manifold invariant of Broda [B1] and (up to an unspecified normalization factor) the state-sum 4-manifold invariant of Crane-Yetter [CY] are equivalent to the signature of the 4-manifold. Subsequently Broda [B2] defined another surgical invariant of 4-manifolds in which the 1- and 2- handles are treated differently. We use a refinement of Roberts' techniques developed in [CKY] to identify the normalization factor to show that the "improved" surgical invariant of Broda [B2] also depends only on the signature and Euler character.

As a starting point, let us first observe that the construction of Crane-Yetter [CY] does not really depend on the use of labels chosen from the irreps of $U_q(sl_2)$ at the principal r^{th} root of unity: the simple objects of any artinian semi-simple tortile category (cf. [S, Y]) in which all objects are self-dual and the fusion rules are multiplicity free will suffice. In particular, if we restrict to the integer spin (bosonic)³ irreps, we obtain a construction of a different invariant of 4-manifolds.

In what follows, we use Temperley-Lieb recoupling theory (cf. [KL,L,R]). In particular, arcs are labelled with elements of $\{0, 1, \dots, r-2\}$ (twice the spin), $A = e^{2\pi i/4r}$, $q = A^2$, $\Delta(n) = (-1)^n \frac{q^{n+1} - q^{-n-1}}{q - q^{-1}}$, $\theta(a, b, c)$ denoted the evaluation of the theta-net with edge labelled a, b , and c , and $15-j$ denotes the evaluation of the Temperley-Lieb version of the Crane-Yetter quantum $15j$ -symbol (with indices suppressed).

We then adopt the following further notational conventions:

Arcs labelled ω denote the linear combination of arcs labelled $0, 1, \dots, r-2$ in which the coefficient of i is $\Delta(i)$. Arcs labelled $\tilde{\omega}$ denote the linear combination of arcs labelled $0, 2, \dots, 2\lfloor \frac{r-2}{2} \rfloor$ (even integers) in which the coefficient of i is $\Delta(i)$. N denotes the sum of the squares of the $\Delta(i)$'s, $\tilde{N}$ denotes the sum of the squares of the $\Delta(i)$'s for i even. Let κ be as in [KL,R], the evaluation of an ω labelled 1-framed unknot divided by the positive square root of N , and let $\tilde{\kappa}$ be the evaluation of an $\tilde{\omega}$ labelled 1-framed unknot divided by $\tilde{N}$.

If L is a framed link, then $\tilde{\omega}(L)$ denotes the evaluation of the link with all components labelled $\tilde{\omega}(L)$. If $\mathcal{L}$ is a set of 4-manifold surgery instructions (cf. Kirby [K]), that is a link L with a

¹Supported by National Science Foundation grant #DMS-9106476

²Supported by National Science Foundation grant #DMS-9205277 and the Program for Mathematics and Molecular Biology of the University of California at Berkeley, Berkeley, CA

³This use of bosonic is a hideous abuse of language—everything in sight has braid statistics—the "bosons" of this paper are the result of q -deforming honest bosons.

distinguished 0-framed unlink $\dot{L}$, then $\mathcal{B}^!(\mathcal{L})$ denotes the evaluation of the link L with all components of $\dot{L}$ (one-handle attachments) colored ω and all other components of L (two-handle attachments) colored $\tilde{\omega}$.

We then have

Lemma 1 *$\tilde{\omega}(L)$ is invariant under handle-sliding. $\mathcal{B}^!\mathcal{L}$ is invariant under handle-sliding of 1- and 2-handles 1-handles and of 2-handles over 2-handles.*

proof: This follows immediately from handle-sliding over components labelled ω and the analysis given in Remark 17 §12.6 of Kauffman/Lins [KL] once it is observed that pairs of bosons only couple to produce bosons. $\square$

And

Lemma 2 (The bosonic encirclement lemma)

$$\sum_{j \text{ even}} \text{diagram} = 0$$

whenever n is even and non-zero.

proof: This follows from the same proof as the encirclement lemma of Lickorish [L] (cf. also Kauffman/Lins [KL]) with the “auxiliary loop” labelled 2 instead of 1. $\square$

Let

$$CY_B(W) = \tilde{N}^{n_0 - n_1} \sum_{\substack{\text{even labellings} \\ \lambda \text{ of faces and} \\ \text{tetrahedra}}} \prod_{\text{faces } \sigma} \Delta(\lambda(\sigma)) \prod_{\substack{\text{tetrahedra} \\ \tau}} \frac{\Delta(\lambda(\sigma))}{\theta(\lambda(\tau), \lambda(\tau_0), \lambda(\tau_2)) \theta(\lambda(\tau), \lambda(\tau_1), \lambda(\tau_3))} \prod_{\text{4-simplexes}} 15 - j$$

be the bosonic Crane-Yetter invariant.

Let $|L|$ (resp. $\nu(L)$, $\sigma(L)$) denote the number of components of a link L (resp. the nullity of the linking matrix of L , the signature of the linking matrix of L).

We can then define a purely bosonic version of Broda’s original invariant by

$$Br_B(W) = \frac{\tilde{\omega}(L)}{\tilde{N}^{\frac{|L| + \nu(L)}{2}}}$$

where L is the underlying link of a surgery presentation of W ; while a bosonic version of the Reshetikhin/Turaev [RT] 3-manifold invariant is given by

$$I_B(M) = \tilde{\kappa}^{-\sigma(L)} \tilde{N}^{\frac{|L| + 1}{2}} \tilde{\omega}(L)$$

where L is a framed link giving surgery instructions for M .

Applying the two lemmas above in an analysis otherwise identical to that of given by Roberts [R] of the original Broda invariant [B1] shows that

Proposition 3

$$Br_B(W) = \tilde{\kappa}^{\sigma(W)}$$

Similarly it follows from the bosonic encirclement lemma that

$$CY_B(W) = \tilde{N}^{n_0-n_1-n_3} \tilde{\omega}(L)$$

where n_d is the number of d -simplexes in a triangulation, and L is the link derived from a triangulation by putting a 0-framed unknot in each tetrahedron, and a loop around each 2-simplex (running mostly through 4-simplexes but linking each tetrahedron's unknot) after the manner of Roberts [R].

It then follows as in [CKY] that

Proposition 4

$$CY_B(W) = \tilde{\kappa}^{\sigma(W)} \tilde{N}^{\frac{\chi(W)}{2}} \quad (*)$$

Now, Broda's new invariant is defined by

$$\mathcal{B}(W) = \frac{\mathcal{B}^!(L)}{\tilde{N}^{\nu(L)} N^{\frac{|L|-\nu(L)}{2}}}$$

For convenience we first analyse a slightly different normalization (for which the proof of invariance is effectively identical to that for $\mathbf{B}(W)$): let

$$\mathbf{B}(W) = \frac{\mathcal{B}^!(L)}{\tilde{N}^{|L-\dot{L}|-|\dot{L}|} N^{|\dot{L}|}}$$

Now, it follows from the original encirclement lemma of Lickorish [L] that

$$CY_B(W) = \tilde{N}^{n_0-n_1} N^{-n_3} \mathcal{B}^!(\mathcal{L}) \quad (**)$$

where $\mathcal{L}$ is the surgery instructions given by associating the link L to the triangulation as above, and letting $\dot{L}$ be the unlink of loops in the tetrahedra.

Observe that $\mathbf{B}$ is multiplicative under connected sum, and that $\mathbf{B}(S^1 \times S^3) = \tilde{N}$ (an easy calculation). As shown in Roberts [R], $\mathcal{L}$ is a surgery presentation for $W \# (\#^{n_4-1} S^1 \times S^3)$.

From this and the fact that for $\mathcal{L}$, $|L - \dot{L}| = n_2$ and $|\dot{L}| = n_3$, we see that

$$\begin{aligned} \frac{\mathcal{B}^!(\mathcal{L})}{\tilde{N}^{n_2-n_3} N^{n_3}} &= \mathbf{B}(W \# (\#^{n_4-1} S^1 \times S^3)) \\ &= \mathbf{B}(W) \tilde{N}^{n_4-1}. \end{aligned}$$

Thus

$$\mathcal{B}^!(L) = \mathbf{B}(W) \tilde{N}^{n_2-n_3+n_4-1} N^{n_3}. \quad (***)$$

It then follows from (*), (**) and (***) that

$$\mathbf{B}(W) = \tilde{\kappa}^{\sigma(W)} \tilde{N}^{\frac{\chi(W)}{2}-1}$$

To return to Broda's [B2] original normalization, note that

$$\mathcal{B}(W) = \mathbf{B}(W) (\tilde{N} N^{-\frac{1}{2}})^{|L-\dot{L}|-|\dot{L}|-\nu(L)}$$

From which we obtain

Theorem 5 *If W is a connected closed oriented smooth 4-manifold, then*

$$\mathcal{B}(W) = \tilde{\kappa}^{\sigma(W)} \left(\frac{\tilde{N}}{N} \right)^{\frac{\chi(W)}{2} - 1}$$

proof: It suffices to shown that if W is given by the surgery instruction $\mathcal{L}$, then

$$|L - \dot{L}| - |\dot{L}| - \nu(L) = \chi(W) - 2.$$

But this follows immediately from the observation that $\nu(L)$ is the number of 3-handles attached in completing the construction of W . $\square$

References

- [B1] Broda, B., *Surgical invariants of 4-manifolds*, preprint (1993).
- [B2] Broda, B., *A Surgical invariant of 4-manifolds*, Proceedings of the Conference on Quantum Topology (D.N. Yetter, ed.), World Scientific, to appear.
- [CKY] Crane, L., Kauffman, L. H. and Yetter, D. N., *Evaluating the Crane-Yetter invariant*, e-preprint hep-th/9309063, and to appear *Quantum Topology* (R. Baadhio and L.H. Kauffman eds.), World Scientific.
- [CY] Crane, L. and Yetter, D. N., *A categorical construction of 4D topological quantum field theories*, e-preprint hep-th/9301062, and to appear in *Quantum Topology* (R. Baadhio and L.H. Kauffman eds.), World Scientific.
- [KL] Kauffman, L. H. and Lins, S. L. *Temperley-Lieb Recoupling Theory and Invariants of 3-Manifolds*, Princeton University Press, to appear.
- [K] Kirby, R., *The topology of 4-manifolds*, SLNM vol. 1374, Springer-Verlag (1989).
- [L] Lickorish, W. B. R. *The Temperley-Lieb algebra and 3-Manifold invariants*, preprint (1992).
- [RT] Reshetikhin, N. and Turaev, V. G., *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. **103** (1991), 547-597.
- [R] Roberts, J., *Skein theory and Turaev-Viro invariants*, preprint (1993).
- [S] Shum, M.-C., *Tortile tensor categories*, Doctoral Dissertation, Macquarie University, 1989.
- [Y] Yetter, D.N., *State-sum invariants of 3-manifolds associated to artinian semisimple tortile categories*, Topology and its App., to appear.